

New York Family Practice Test Requisition Form

Ordering Physician/Laboratory

(Required: Include the ordering physician's first & last name, NPI, practice name, complete address, phone number and fax number.)

Patient Information (Please Print)

Name (Last, First) (Required):

In Care of:

Patient Address:

City: _____ State: _____ Zip: _____

Assigned Sex at Birth (Required): _____ Date of Birth (Required): _____ Patient ID#:

Female Male

Phone Number (Required): _____

Cell Phone
 Home Phone

Race: Alaska Native or American Indian Asian Black or African American Multiracial Native Hawaiian or other Pacific Islander Other race White Does not wish to disclose Not provided

Ethnicity: Hispanic or Latino
 Not Hispanic or Latino
 Unknown

Physician to receive additional result report:

Physician's Signature: _____ Date: _____

Specimen Information

Date Collected (Required): Specimen Type: Specimen Source:

OneSwab®: _____
 Blood
 Saliva
 UroSwab®

Was **UroSwab®** collected from a catheterized patient? Yes

Sexually Transmitted Infections - OneSwab®

Common ICD10 codes (required):

N89.8 Other specified noninflammatory disorders of vagina
 Z20.2 Contact with and (suspected) exposure to infections with a predominantly sexual mode of transmission
 N76.4 Abscess of vulva
 Other: _____

121 **Leukorrhea Panel** Includes -
 105 *Chlamydia trachomatis* (*Reflex to antibiotic resistance by Molecular Analysis),
 167 *Neisseria gonorrhoeae*,
 111 *Trichomonas vaginalis* (*Reflex to **M** resistance),
 129 *Mycoplasma genitalium* (*Reflex to antibiotic resistance by Molecular Analysis)

115 **Genital Ulcer Disease Panel** Includes -
 122 *Haemophilus ducreyi* 126 Herpes subtype (HSV-1, HSV-2) 110 *Treponema pallidum* (syphilis)

739 **HPV Type-Detect® 2.0 by Bio-plex Analysis (High Risk Subtypes Only)**

Vaginitis & Vaginosis - OneSwab®

Common ICD10 codes (required):

N76.0 Acute vaginitis
 N89.8 Other specified noninflammatory disorders of vagina
 R10.2 Pelvic and perineal pain
 Other: _____

759 **Bacterial Vaginosis (BV) Panel with Lactobacillus Profiling by qPCR** Includes -
Famyhessea vaginæ (*Atopobium vaginæ*), BVAB1, BVAB2, BVAB3, *Bacteroides fragilis*, *Bifidobacterium breve*, *Megasphaera* Type 1 & 2, *Gardnerella vaginalis*, *Mobiluncus curtisi*, *M. mulieris*, *Prevotella bivia*, *Sneathia sanguinegens*, *Streptococcus anginosus*

182 **Aerobic Vaginitis (AV) Panel** Includes -
 127 Group B Streptococcus (GBS) 141 *Escherichia coli*
 184 *Staphylococcus aureus* 153 *Enterococcus faecalis*

6009 **Candida Panel Basic with fluconazole resistance by X-Plate Technology®** Includes -
 551 *Candida albicans* 559 *Candida glabrata*
 558 *Candida parapsilosis* 557 *Candida tropicalis*
 566 *Candida krusei*

134 **Urogenital Mycoplasma & Ureaplasma Panel**
 129 *Mycoplasma genitalium* 130 *Mycoplasma hominis*
 320 *Ureaplasma urealyticum*

Intestinal Pathogens - OneSwab®

Loose Stool, Rectal Swab, No Lubricant

Common ICD10 codes (required):

R19.7 Diarrhea, unspecified
 R10.9 Unspecified abdominal pain

R11.2 Nausea with vomiting, unspecified
 Other: _____

365 *Campylobacter jejuni*
 162 *Clostridium difficile* (Toxins A and B)
 168 *Escherichia coli* (O157 and Shiga toxin)

370 *Giardia intestinalis*
 310 *Helicobacter pylori*
 158 *Listeria monocytogenes*
 160 *Salmonella*
 161 *Shigella*

Other Tests/Panels:

ICD10 codes (required):

For a full menu of testing, please visit www.mdlab.com

Skin & Soft Tissue Infections - OneSwab®

Common ICD10 codes (required):

A49.9 Bacterial infection, unspecified B99.8 Other infectious disease
 B99.9 Unspecified infectious disease (opportunistic) Other:
373 **Skin & Soft Tissue Infections (STTI) Panel (NY Only)** Includes -
 153 *Enterococcus faecalis* 141 *Escherichia coli* 1112 Group A Streptococcus
 127 Group B Streptococcus (GBS) 172 *Klebsiella Species*
 1118 MRSA: Methicillin Resistant and Methicillin Susceptible (MSSA) *Staphylococcus aureus* by Conventional PCR 1119 CA-MRSA: Community-Associated MRSA. Panton-Valentine Leukocidin (PVL) DNA** (Type IV MRSA + #1118 Req.) [Community Associated MRSA = Type IV MRSA+ and PVL+] 146 *Proteus mirabilis* 174 *Pseudomonas aeruginosa*

367 **STTI Panel Antibiotic Resistance** Includes - *[E. faecalis, E. coli, GAS, GBS, K. oxytoca, K. pneumoniae, P. mirabilis, P. aeruginosa, MRSA: AC, AM (for E. faecalis), C, CL, D, TS, CP, CF, PT, I, G]* (153, 141, 1112, 127, 172, 146, 174, or 1118 Req.) When panel is ordered and individual tests are not selected, all 8 will be performed & billed)

UroSwab®

Common ICD10 codes (required):

Z20.2 Contact with and (suspected) exposure to infections with a predominantly sexual mode of transmission R30.9 Painful micturition, unspecified
 R36.9 Urethral discharge, unspecified Other: _____

SEXUALLY TRANSMITTED INFECTIONS

Applicable for adolescent females who are not candidates for pelvic exams.

121 Leukorrhea Panel

Includes -
 105 *Chlamydia trachomatis* (*Reflex to antibiotic resistance by Molecular Analysis)
 167 *Neisseria gonorrhoeae*
 111 *Trichomonas vaginalis* (*Reflex to **M** resistance)
 129 *Mycoplasma genitalium* (*Reflex to antibiotic resistance by Molecular Analysis)

109 *N. gonorrhoeae & C. trachomatis**

Common ICD10 codes (required):

N39.0 Urinary tract infection, site not specified R30.1 Vesical tenesmus
 R30.0 Dysuria Other: _____

URINARY TRACT INFECTIONS

369 *Acinetobacter baumannii* complex 578 *Candida kefyr*
 149 *Actinomyces* spp. group 1 577 *Candida lusitaniae*
 576 *Candida dubliniensis* 574 *Candida utilis*

Pharmacogenomics- whole blood or saliva

ICD10 codes (required):

Note: This testing requires a signed Patient Informed Consent/Insurance Acknowledgment Form

3407 **Depressive Disorder & Major Depressive Disorder** - Amitriptyline, Antidepressants (gen), Antipsychotics (gen), Aripiprazole, Bupropion, Citalopram, Clomipramine, Desipramine, Desvenlafaxine, Diazepam, Doxepin, Duloxetine, Escitalopram, Fluoxetine, Fluvoxamine, Imipramine, Maprotiline, Milnacipran, Mirtazapine, Nortriptyline, Olanzapine, Opipramol, Paroxetine, Quetiapine, Sertraline, SSRIs (gen), Trimipramine, Venlafaxine, Vilazodone, Vortioxetine (ABC1, ADRA2A, ANKK1, COMT, CYP1A2, CYP2B6, CYP2C19, CYP2C9, CYP2D6, CYP3A4, CYP3A5, DRD4, GABRP, GRIK4, HTR2A, HTR2C, MTHFR, UGT1A1)

3404 **Anxiety, Insomnia, Severe Agitation** - Alprazolam, Bupropion, Daridorexant, Desvenlafaxine, Dexmedetomidine, Duloxetine, Escitalopram, Lemborexant, Lorazepam, Midazolam, Oxazepam, Pimavanserin, Valbenazine, Venlafaxine, Vilazodone (ABC1, ADRA2A, ANKK1, COMT, CYP1A2, CYP2B6, CYP2C19, CYP2D6, CYP3A4, GABRP, HTR2A, UGT1A1, UGT2B15)

3406 **Bipolar Disorder** - Aripiprazole, Asenapine, Cariprazine, Clonazepam, Lamotrigine, Lithium, Lumateperone, Lurasidone, Olanzapine, Ocarbazepine, Quetiapine, Samidorphan, Valbenazine, Valproic acid, Vilazodone (ABC1, ABC2, ANKK1, CYP1A2, CYP2C9, CYP2D6, CYP3A4, CYP3A5, COMT, DRD1, HTR2A, MTHFR, UGT1A1)

3402 **ADD/ADHD** - Atomoxetine, Bupropion, Dextroamphetamine, Imipramine, Lisdexamfetamine, Methylphenidate, Modafinil, Nortriptyline, Viloxazine (ANKK1, ABC1, ADRA2A, CYP2B6, CYP2C19, CYP2D6, DRD1)

Confirmation of Informed Consent and Medical Necessity for Pharmacogenomic/Genetic Testing
 My signature below acknowledges the patient has been informed about the purpose, limitation and possible risks of genetic testing. The patient has been given the opportunity to ask questions about this consent and seek outside genetic counseling.

If the genetic testing is covered by the patient's health plan and the out-of-pocket expense is less than \$150.00, testing will proceed without further delay or additional contact. The patient's signed informed consent is being provided with this requisition. I confirm that this testing is medically necessary for the specified patient and that these results will be used in the medical management and treatment decisions for this patient.

Medical Professional Signature (Req.): _____ Date: _____

Refer to the back for antibiotic abbreviation key.

IH0320 Rel.: 2.12.2026

Antibiotic Abbreviations Key

A = aztreonam **AC**= amoxicillin-clavulanic acid, **AP** = ampicillin, **AZ** = azithromycin, **CC** = ceftriaxone/ceftriaxime, **C** = cephalothin (cephalexin), **CF** = cefepime, **CP** = ciprofloxacin, **CL** = clindamycin, **D** = doxycycline, **F** = fosfomycin, **FL** = fluoroquinolone **G** = gentamicin, **I** = imipenem, **L** = linezolid, **M** = metronidazole **N** = nitrofurantoin, **PT** = piperacillin-tazobactam, **TS** = trimethoprim-sulfamethoxazole.

Medical Necessity Guidelines:

Physicians must only order tests that they have determined are medically necessary for the diagnosis and treatment of a patient. MDL offers individual tests, as well as a limited number of customized panels. MDL provides practitioners with the flexibility to choose appropriate individual tests for each specimen to assure that the convenience of ordering panels does not impede them from ordering tests/panels that are medically necessary. All tests listed in panels may be ordered individually using this test requisition form. If you choose to order a panel, please make certain that each and every test is medically necessary. If you check off a panel as your choice, MDL understands that the physician has determined that all of the component tests are medically necessary, and will perform, report and bill for all such component tests.

** This test can only be performed when the test in parenthesis is positive. All tests performed will be billed.
Test by Real-Time PCR unless otherwise specified.

OneSwab® & UroSwab® are registered in the USPTO.

Specimen Collection Platform		TAT*	Stability	Test Additions [¥]	Specimen Collection
OneSwab®		24 - 72 hours	7 days	30 days to add tests	1. Collect specimen with the sterile swab provided. 2. Insert swab into the transport media, break off swab handle and tightly re-secure the cap on the transport media vial.
UroSwab®		24 - 72 hours	4 days	14 days to add tests	1. Have patient collect a urine specimen in a collection cup. 2. Dip the sponge swab into collection cup to absorb the urine. 3. Tightly re-secure the cap on the vial.
Blood	Yellow top tube (ACD solution A)	3 - 5 days	48 Hours	30 days to add tests	1. In accordance with the standard operating procedure of your facility, collect blood in two yellow top (ACD solution A) tubes. 2. Allow the tubes to fill properly to ensure the proper blood to anticoagulant ratio. 3. Invert gently several times to mix and prevent clot formation. Do not shake the tubes. Do not centrifuge.
Saliva		5 - 10 days	48 Hours	--	<ul style="list-style-type: none"> • Vigorously rinse your mouth with clean water 5 minutes prior to specimen collection (30 minutes prior is ideal). • After rinsing, do not brush your teeth, use mouthwash, eat, drink, chew gum or smoke prior to sample collection. 1. Begin collecting your sample by allowing saliva to pool in your mouth. Then spit into the wide funnel of the tube allowing saliva to collect in the upper chamber of the tube. Fill the tube until the amount of saliva (not bubbles) reaches the fill line as shown. 2. Once filled, unscrew the funnel allowing the saliva to flow into the lower chamber of the tube containing the stabilizing solution. Discard the funnel. 3. Use the blue cap to close the tube tightly. Shake the capped tube for 5 seconds.

* Up to 72 hours with reflex/antibiotic resistance testing

[¥] Pending QC review for sufficient specimen volume

Specimen Packaging:

1. Label every vial with a minimum of 2 patient identifiers including the patient's name and date of birth. Be sure the name on the vial is written exactly the same way as on the test requisition form.
2. Place the vial into the Styrofoam/Cardboard container. You can fit up to 3 vials in one container.
3. Place the Styrofoam/Cardboard container into the central pocket of a biohazard bag containing absorbent material.
4. Place a completed test requisition form for each vial in the front pocket of the biohazard bag.
5. Place the biohazard bag into the prepaid Lab pack Envelope that has a preaddressed airbill attached. One envelope will accommodate 6-7 containers. Package as many containers in one Labpack as possible.
6. Be sure to seal the Lab pack by removing the plastic from the top of the adhesive.

Specimen Pick-up:

- If you have a specimen pick-up for a local courier in the NJ, PA, DE, D.C., MD, VA, KS, MO or Phoenix, AZ area, please call (877) 205-0005 no later than 2 hours prior to the closing of your facility.
- If you have a specimen pick-up, please call your sales representative or MDL customer service at 877.269.0090 no later than 2 hours prior to the closing of your facility.

Helpful Hints Checklist

Please review these helpful hints to reduce specimen discrepancies and enhance turnaround time.

Verify Patient Name - did you:

- ✓ attach the correct demographics sheet?
- ✓ write the patient's name on the requisition form?

Patient Name Matches on Vial & Requisition Form- did you:

- ✓ make sure names on vial and requisition form match?
- ✓ list the patients married or maiden name?
- ✓ list a nickname by mistake?

Verify Date of Collection- did you:

- ✓ write the correct year?
- ✓ write the correct month?
- ✓ list the date of birth instead?

Verify Tests- did you:

- ✓ clearly mark each box?
- ✓ order tests appropriate for the specimen type?

No Tests Ordered- did you:

- ✓ mark the boxes for the tests/panels ordered?

Supply Orders:

Easily place supply orders online by visiting our website:



<http://www.mdlab.com/physicians/supplies/#>

Supply orders may also be placed by contacting our Client Services department toll free 877.269.0090 or by fax 609.570.1050. Supply requests are processed and shipped on a daily basis. Please allow 3 to 5 business days for delivery, depending upon your location.

MDL Contact Information



GBS Hotline

24 hours - 7 days a week
Group B Strep & HSV results only

877.MDL.GBS7

877.635.4277

Quality Control Department For Technical Assistance

877.269.0090

609.245.7665

Client Services

General Questions, Results

877.269.0090

609.570.1050

Client Services

Billing Questions

877.333.9233

609.245.7683